

Distributed multi-agent optimization with state-dependent communication

Ilan Lobel · Asuman Ozdaglar · Diego Feijer

Received: 4 April 2010 / Accepted: 9 January 2011 / Published online: 4 June 2011
© Springer and Mathematical Optimization Society 2011

Abstract We study distributed algorithms for solving global optimization problems in which the objective function is the sum of local objective functions of agents and the constraint set is given by the intersection of local constraint sets of agents. We assume that each agent knows only his own local objective function and constraint set, and exchanges information with the other agents over a randomly varying network topology to update his information state. We assume a *state-dependent communication model* over this topology: communication is Markovian with respect to the states of the agents and the probability with which the links are available depends on the states of the agents. We study a *projected multi-agent subgradient algorithm* under state-dependent communication. The state-dependence of the communication introduces significant challenges and couples the study of information exchange with the

This paper is dedicated to the memory of Paul Tseng, a great researcher and friend.

This research was partially supported by the National Science Foundation under Career grant DMI-0545910, the DARPA ITMANET program, and the AFOSR MURI R6756-G2.

Electronic supplementary material The online version of this article (doi:[10.1007/s10107-011-0467-x](https://doi.org/10.1007/s10107-011-0467-x)) contains supplementary material, which is available to authorized users.

I. Lobel (✉)
Stern School of Business, New York University, New York, NY, USA
e-mail: ilobel@stern.nyu.edu

A. Ozdaglar · D. Feijer
Laboratory for Information and Decision Systems, Department of Electrical Engineering and Computer Science, Massachusetts Institute of Technology, Cambridge, MA, USA
e-mail: asuman@mit.edu

D. Feijer
e-mail: feijer@mit.edu

analysis of subgradient steps and projection errors. We first show that the multi-agent subgradient algorithm when used with a constant stepsize may result in the agent estimates to diverge with probability one. Under some assumptions on the stepsize sequence, we provide convergence rate bounds on a “disagreement metric” between the agent estimates. Our bounds are time-nonhomogeneous in the sense that they depend on the initial starting time. Despite this, we show that agent estimates reach an almost sure consensus and converge to the same optimal solution of the global optimization problem with probability one under different assumptions on the local constraint sets and the stepsize sequence.

Mathematics Subject Classification (2000) 90C30-Nonlinear Programming · 90B15-Network Models, Stochastic

1 Introduction

Due to computation, communication, and energy constraints, several control and sensing tasks are currently performed collectively by a large network of autonomous agents. Applications are vast including a set of sensors collecting and processing information about a time-varying spatial field (e.g., to monitor temperature levels or chemical concentrations), a collection of mobile robots performing dynamic tasks spread over a region, mobile relays providing wireless communication services, and a set of humans aggregating information and forming beliefs about social issues over a network. These problems motivated a large literature focusing on design of optimization, control, and learning methods that can operate using local information and are robust to dynamic changes in the network topology. The standard approach in this literature involves considering “consensus-based” schemes, in which agents exchange their local estimates (or states) with their neighbors with the goal of aggregating information over an *exogenous* (fixed or time-varying) network topology. In many of the applications, however, the relevant network topology is configured *endogenously as a function of the agent states*, for example, the communication network varies as the location of mobile robots changes in response to the objective they are trying to achieve. A related set of problems arises when the current information of decentralized agents influences their potential communication pattern, which is relevant in the context of sensing applications and in social settings where disagreement between the agents would put constraints on the amount of communication among them.

In this paper, we propose a general framework for design and analysis of distributed multi-agent optimization algorithms with state dependent communication. Our model involves a network of m agents, each endowed with a local objective function $f_i : \mathbb{R}^n \rightarrow \mathbb{R}$ and a local constraint $X_i \subseteq \mathbb{R}^n$ that are private information, i.e., each agent only knows its own objective and constraint. The goal is to design distributed algorithms for solving a global constrained optimization problem for optimizing an objective function, which is the sum of the local agent objective functions, subject to a constraint set given by the intersection of the local constraint sets of the agents. These algorithms involve each agent maintaining an estimate (or state) about the optimal

solution of the global optimization problem and update this estimate based on local information and processing, and information obtained from the other agents.

We assume that agents communicate over a network with randomly varying topology. Our random network topology model has two novel features: First, we assume that the communication at each time instant k , (represented by a *communication matrix* $A(k)$ with positive entries denoting the availability of the links between agents) is Markovian on the states of the agents. This captures the time correlation of communication patterns among the agents.¹ The second, more significant feature of our model is that the probability of communication between any two agents at any time is a function of the agents' states, i.e., the closer the states of the two agents, the more likely they are to communicate. As outlined above, this feature is essential in problems where the state represents the position of the agents in sensing and coordination applications. For example, consider a prospecting application, where our objective is to select a drilling location using a family of mobile robots. Each robot is potentially outfitted with a different sensor – some robots are capable of detecting the desired minerals underground, while others are used to determine how difficult it would be to drill in each potential location. The desired outcome is to have the mobile robots converge on the optimal drilling location, despite the fact that the further away the robots are from each other, the less likely it is that they are able to communicate with each other.

For this problem, we study a *projected multi-agent subgradient algorithm*, which involves each agent performing a local averaging to combine his estimate with the other agents' estimates he has access to, taking a subgradient step along his local objective function, and projecting the estimates on his local constraint set. We represent these iterations as stochastic linear time-varying update rules that involve the agent estimates, subgradients and projection errors explicitly. With this representation, the evolution of the estimates can be written in terms of stochastic transition matrices $\Phi(k, s)$ for $k \geq s \geq 0$, which are products of communication matrices $A(t)$ over a window from time s to time k . The transition matrices $\Phi(k, s)$ represent aggregation of information over the network as a result of local exchanges among the agents, i.e., in the long run, it is desirable for the transition matrices to converge to a uniform distribution, hence aligning the estimates of the agents with uniform weights given to each (ensuring that information of each agent affects the resulting estimate uniformly). As a result, the analysis of our algorithm involves studying convergence properties of transition matrices, understanding the limiting behavior of projection errors, and finally studying the algorithm as an "approximate subgradient algorithm" with bounds on errors due to averaging and projections.

In view of the dependence of information exchange on the agent estimates, it is not possible to decouple the effect of stepsizes and subgradients from the convergence of the transition matrices. We illustrate this point by first presenting an example in which the projected multi-agent subgradient algorithm is used with a constant stepsize $\alpha(k) = \alpha$ for all $k \geq 0$. We show that in this case, agent estimates and the

¹ Note that our model can easily be extended to model Markovian dependence on other stochastic processes, such as channel states, to capture time correlation due to global network effects. We do not do so here for notational simplicity.

corresponding global objective function values may diverge with probability one for any constant value of the stepsize. This is in contrast to the analysis of multi-agent algorithms over exogenously varying network topologies where it is possible to provide error bounds on the difference between the limiting objective function values of agent estimates and the optimal value as a function of the constant stepsize α (see [14]).

We next adopt an assumption on the stepsize sequence $\{\alpha(k)\}$ (see Assumption 5), which ensures that $\alpha(k)$ decreases to zero sufficiently fast, while satisfying $\sum_{k=0}^{\infty} \alpha(k) = \infty$ and $\sum_{k=0}^{\infty} \alpha^2(k) < \infty$ conditions. Under this assumption, we provide a bound on the expected value of the disagreement metric, defined as the difference $\max_{i,j} |\Phi(k, s)_{ij} - \frac{1}{m}|$. Our analysis is novel and involves constructing and bounding (uniformly) the probability of a hierarchy of events, the length of which is specifically tailored to grow faster than the stepsize sequence, to ensure propagation of information across the network before the states drift away too much from each other. In contrast to exogenous communication models, our bound is *time-nonhomogeneous*, i.e., it depends on the initial starting time s as well as the time difference $(k - s)$. We also consider the case where we have the assumption that the agent constraint sets X_i 's are compact, in which case we can provide a bound on the disagreement metric without any assumptions on the stepsize sequence.

Our next set of results study the convergence behavior of agent estimates under different conditions on the constraint sets and stepsize sequences. We first study the case when the local constraint sets of agents are the same, i.e., for all i , $X_i = X$ for some nonempty closed convex set. In this case, using the time-nonhomogeneous contraction provided on the disagreement metric, we show that agent estimates reach almost sure consensus under the assumption that stepsize sequence $\{\alpha(k)\}$ converges to 0 sufficiently fast (as stated in Assumption 5). Moreover, we show that under the additional assumption $\sum_{k=0}^{\infty} \alpha(k) = \infty$, the estimates converge to the same optimal point of the global optimization problem with probability one. We then consider the case when the constraint sets of the agents X_i are different convex compact sets and present convergence results both in terms of almost sure consensus of agent estimates and almost sure convergence of the agent estimates to an optimal solution under weaker assumptions on the stepsize sequence.

Our paper contributes to the growing literature on multi-agent optimization, control, and learning in large-scale networked systems. Most work in this area builds on the seminal work by Tsitsiklis et al. [25] and Bertsekas and Tsitsiklis [3], which developed a general framework for parallel and distributed computation among different processors. Our work is related to different strands of literature in this area. One strand focuses on reaching consensus on a particular scalar value or computing exact averages of the initial values of the agents, as natural models of cooperative behavior in networked-systems (for deterministic models, see [8, 13, 21] and [22]); for randomized models, where the randomness may be due to the choice of the randomized communication protocol or due to the unpredictability in the environment that the information exchange takes place, see [10, 7, 12, 26, 24]. Another recent literature studies optimization of more general objective functions using subgradient algorithms and consensus-type mechanisms (see [16, 14, 17–19, 23, 27]). Of particular relevance to our work are the papers [14] and [19]. In [14], the authors studied a multi-agent

unconstrained optimization algorithm over a random network topology which varies independently over time and established convergence results for diminishing and constant stepsize rules. The paper [19] considered multi-agent optimization algorithms under deterministic assumptions on the network topology and with constraints on agent estimates. It provided a convergence analysis for the case when the agent constraint sets are the same. The paper [9] uses a distributed version of the dual averaging algorithm of [20] in a randomized network model and obtains bounds on the convergence rate as a function of the network topology. A related, but somewhat distinct literature, uses consensus-type schemes to model opinion dynamics over social networks (see [1,4,5,11]). Among these papers, most related to our work are [4] and [5], which studied dynamics with opinion-dependent communication, but without any optimization objective.

The rest of the paper is organized as follows: in Sect. 2, we present the optimization problem, the projected subgradient algorithm and the communication model. In Sect. 3, we introduce and bound the disagreement metric ρ , which determines the spread of information in the network. Section 4 shows the convergence of the projected subgradient methods and Sect. 5 concludes. Appendix A includes some notation and basic relations, online Appendix B contains omitted proofs and online Appendix C contains a simulation analysis. Appendices B and C are available online [15].

2 The model

2.1 Optimization model

We consider a network that consists of a set of nodes (or agents) $\mathcal{M} = \{1, \dots, m\}$. We assume that each agent i is endowed with a local objective (cost) function f_i and a local constraint function X_i and this information is distributed among the agents, i.e., each agent knows only his own cost and constraint component. Our objective is to develop distributed algorithms that can be used by these agents to cooperatively solve the following constrained optimization problem:

$$\begin{aligned} & \text{minimize} && \sum_{i=1}^m f_i(x) && (1) \\ & \text{subject to} && x \in \bigcap_{i=1}^m X_i, \end{aligned}$$

where each $f_i : \mathbb{R}^n \rightarrow \mathbb{R}$ is a convex (not necessarily differentiable) function, and each $X_i \subseteq \mathbb{R}^n$ is a closed convex set. We denote the intersection set by $X = \bigcap_{i=1}^m X_i$ and assume that it is nonempty throughout the paper. Let f denote the global objective, that is, $f(x) = \sum_{i=1}^m f_i(x)$, and f^* denote the optimal value of problem (1), which we assume to be finite. We also use $X^* = \{x \in X : f(x) = f^*\}$ to denote the set of optimal solutions and assume throughout that it is nonempty.

We study a distributed multi-agent subgradient method, in which each agent i maintains an *estimate* of the optimal solution of problem (1) (which we also refer to as the *state of agent i*), and updates it based on his local information and information

exchanged with other neighboring agents. Every agent i starts with some initial estimate $x_i(0) \in X_i$. At each time k , agent i updates its estimate according to the following:

$$x_i(k + 1) = P_{X_i} \left[\sum_{j=1}^m a_{ij}(k)x_j(k) - \alpha(k)d_i(k) \right], \tag{2}$$

where P_{X_i} denotes the projection on agent i constraint set X_i , the vector $[a_{ij}(k)]_{j \in \mathcal{M}}$ is a vector of weights for agent i , the scalar $\alpha(k) > 0$ is the stepsize at time k , and the vector $d_i(k)$ is a subgradient of agent i objective function $f_i(x)$ at his estimate $v_i(k) = \sum_{j=1}^m a_{ij}(k)x_j(k)$. Hence, in order to generate a new estimate, each agent combines the most recent information received from other agents with a step along the subgradient of its own objective function, and projects the resulting vector on its constraint set to maintain feasibility. We refer to this algorithm as the *projected multi-agent subgradient algorithm*.² Note that when the objective functions f_i are identically zero and the constraint sets $X_i = \mathbb{R}^n$ for all $i \in \mathcal{M}$, then the update rule (2) reduces to the classical averaging algorithm for *consensus* or *agreement* problems, as studied in [6] and [13].

In the analysis of this algorithm, it is convenient to separate the effects of different operations used in generating the new estimate in the update rule (2). In particular, we rewrite the relation in Eq. (2) equivalently as follows:

$$v_i(k) = \sum_{j=1}^m a_{ij}(k)x_j(k), \tag{3}$$

$$x_i(k + 1) = v_i(k) - \alpha(k)d_i(k) + e_i(k), \tag{4}$$

$$e_i(k) = P_{X_i}[v_i(k) - \alpha(k)d_i(k)] - (v_i(k) - \alpha(k)d_i(k)). \tag{5}$$

This decomposition allows us to generate the new estimate using a *linear update rule* in terms of the other agents’ estimates, the subgradient step, and the projection error e_i . Hence, the nonlinear effect of the projection operation is represented by the projection error vector e_i , which can be viewed as a perturbation of the subgradient step of the algorithm. In the sequel, we will show that under some assumptions on the agent weight vectors and the subgradients, we can provide upper bounds on the projection errors as a function of the stepsize sequence, which enables us to study the update rule (2) as an approximate subgradient method.

We adopt the following standard assumption on the subgradients of the local objective functions f_i .

Assumption 1 (*Bounded Subgradients*) The subgradients of each of the f_i are uniformly bounded, i.e., there exists a scalar $L > 0$ such that for every $i \in \mathcal{M}$ and any $x \in \mathbb{R}^n$, we have $\|d\| \leq L$ for all $d \in \partial f_i(x)$.

² See also [19] where this algorithm is studied under deterministic assumptions on the information exchange model and the special case $X_i = X$ for all i .

2.2 Network communication model

We define the *communication matrix* for the network at time k as $A(k) = [a_{ij}(k)]_{i,j \in \mathcal{M}}$. We assume a probabilistic communication model, in which the sequence of communication matrices $A(k)$ is assumed to be Markovian on the *state variable* $x(k) = [x_i(k)]_{i \in \mathcal{M}} \in \mathbb{R}^{n \times m}$. Formally, let $\{n(k)\}_{k \in \mathbb{N}}$ be an independent sequence of random variables defined in a probability space $(\Omega, \mathcal{F}, P) = \prod_{k=0}^{\infty} (\Omega', \mathcal{F}', P')_k$, where $\{(\Omega', \mathcal{F}', P')_k\}_{k \in \mathbb{N}}$ constitutes a sequence of identical probability spaces. We assume there exists a function $\psi : \mathbb{R}^{n \times m} \times \Omega' \rightarrow \mathbb{R}^{m \times m}$ such that $A(k) = \psi(x(k), n(k))$. This Markovian communication model enables us to capture settings where the agents' ability to communicate with each other depends on their current estimates.

We assume there exists some underlying communication graph $(\mathcal{M}, \mathcal{E})$ that represents a 'backbone' of the network. That is, for each edge $e \in \mathcal{E}$, the two agents linked by e systematically attempt to communicate with each other [see Eq. (6) for the precise statement]. We do not make assumptions on the communication (or lack thereof) between agents that are not adjacent in $(\mathcal{M}, \mathcal{E})$. We make the following connectivity assumption on the graph $(\mathcal{M}, \mathcal{E})$.

Assumption 2 (Connectivity) The graph $(\mathcal{M}, \mathcal{E})$ is strongly connected.

The central feature of the model introduced in this paper is that the probability of communication between two agents is potentially small if their estimates are far apart. We formalize this notion as follows: for all $(j, i) \in \mathcal{E}$, all $k \geq 0$ and all $\bar{x} \in \mathbb{R}^{m \times n}$,

$$P(a_{ij}(k) \geq \gamma | x(k) = \bar{x}) \geq \min \left\{ \delta, K / \|\bar{x}_i - \bar{x}_j\|^C \right\}, \tag{6}$$

where K and C are real positive constants, and $0 < \delta, \gamma \leq 1$. We included the parameter δ in the model to upper bound the probability of communication when $\|\bar{x}_i - \bar{x}_j\|^C$ is small. This model states that, for any two nodes i and j with an edge between them, if estimates $x_i(k)$ and $x_j(k)$ are close to each other, then there is a probability at least δ that they communicate at time k . However, if the two agents are far apart, the probability they communicate can only be bounded by the inverse of a polynomial of the distance between their estimates $\|x_i(k) - x_j(k)\|$. If the estimates were to represent physical locations of wireless sensors, then this bound would capture fading effects in the communication channel.

We make two more technical assumptions to guarantee, respectively, that the communication between the agents preserves the average of the estimates, and the agents do not discard their own information.

Assumption 3 (Doubly Stochastic Weights) The communication matrix $A(k)$ is doubly stochastic for all $k \geq 0$, i.e., for all $k \geq 0, a_{ij}(k) \geq 0$ for all $i, j \in \mathcal{M}$, and $\sum_{i=1}^m a_{ij}(k) = 1$ for all $j \in \mathcal{M}$ and $\sum_{j=1}^m a_{ij}(k) = 1$ for all $i \in \mathcal{M}$ with probability one.

The doubly stochasticity assumption on the matrices $A(k)$ is satisfied when agents coordinate their weights when exchanging information, so that $a_{ij}(k) = a_{ji}(k)$ for all $i, j \in \mathcal{M}$ and $k \geq 0$.³

Assumption 4 (*Self Confidence*) There exists $\gamma > 0$ such that $a_{ii} \geq \gamma$ for all agents $i \in \mathcal{M}$ with probability one.

The self-confidence assumption states that each agent gives a significant weight to its own estimate. The constant γ in Eq. (6) is the same as the one used in Assumption 4, such that agents always place a significant weight to the estimates of agents they communicate with as well.

2.3 A counterexample

In this subsection, we construct an example to demonstrate that the algorithm defined in Eqs. (3)–(5) does not necessarily solve the optimization problem given in Eq. (1). The following proposition shows that there exist problem instances where Assumptions 1–4 hold and $X_i = X$ for all $i \in \mathcal{M}$, however the sequence of estimates $x_i(k)$ (and the sequence of function values $f(x_i(k))$) diverge for some agent i with probability one.

The proof of this proposition considers a network consisting of two agents solving a one-dimensional minimization problem. The first agent's objective function is $f_1(x) = -x$, while the second agent's objective function is $f_2(x) = 2x$ and both agents' feasible sets are equal to $X_1 = X_2 = [0, \infty)$. We show that under a certain communication model that satisfies Eq. (6), agent 1's estimate $x_1(k)$ diverges to infinity instead of converging to the optimal solution set $X^* = \{0\}$.

Proposition 1 *Let Assumptions 1, 2, 3 and 4 hold and let $X_i = X$ for all $i \in \mathcal{M}$. Let $\{x_i(k)\}$ be the sequences generated by the algorithm (3)–(5). Let $C > 1$ in Eq. (6) and let the stepsize be a constant value α . Then, there does not exist a bound $M(m, L, \alpha) < \infty$ such that $\liminf_{k \rightarrow \infty} |f(x_i(k)) - f^*| \leq M(m, L, \alpha)$ with probability 1, for all agents $i \in \mathcal{M}$.*

The proposition above shows the algorithm given by Eqs. (3)–(5) does not, in general, solve the global optimization problem (1). However, there are two important caveats when considering the implications of this negative result. The first one is that the proposition only applies if $C > 1$. We leave it is an open question whether the same proposition would hold if $C \leq 1$. The second and more important caveat is that we considered using a constant stepsize in Proposition 1. The stepsize is typically a design choice and, thus, could be chosen to be diminishing in k rather than a constant. In the subsequent sections, we prove that the algorithm given by Eqs. (3)–(5) does indeed solve the optimization problem of Eq. (1), under appropriate assumptions on the stepsize sequence.

³ This will be achieved when agents exchange information about their estimates and “planned” weights simultaneously and set their actual weights as the minimum of the planned weights; see [18] where such a coordination scheme is described in detail.

3 Analysis of information exchange

In this section, we consider how some information that a given agent i obtains at time s affects a different agent j 's estimate $x_j(k)$ at a later time $k \geq s$. In particular, we introduce a disagreement metric $\rho(k, s)$ that establishes how far some information obtained by a given agent at time s is from being completely disseminated in the network at time k . The two propositions at the end of this section provide bounds on $\rho(k, s)$ under two different sets of assumptions.

In view of the linear representation in Eqs. (3)–(5), we can express the evolution of the estimates using products of matrices: for any $s \geq 0$ and any $k \geq s$, we define the *transition matrices* as $\Phi(k, s) = A(s)A(s + 1) \cdots A(k - 1)A(k)$ for all s and k with $k \geq s$. Using the transition matrices, we can relate the estimates at time k to the estimates at time $s < k$ as follows: for all i , and all k and s with $k > s$,

$$\begin{aligned}
 x_i(k + 1) &= \sum_{j=1}^m [\Phi(k, s)]_{ij} x_j(s) \\
 &\quad - \sum_{r=s+1}^k \sum_{j=1}^m [\Phi(k, r)]_{ij} \alpha(r - 1) d_j(r - 1) - \alpha(k) d_i(k) \\
 &\quad + \sum_{r=s+1}^k \sum_{j=1}^m [\Phi(k, r)]_{ij} e_j(r - 1) + e_i(k). \tag{7}
 \end{aligned}$$

Observe from the iteration above that $[\Phi(k, s)]_{ij}$ determines how the information agent i obtains at period $s - 1$ impacts agent j 's estimate at period $k + 1$. If $[\Phi(k, s)]_{ij} = 1/m$ for all agents j , then the information agent i obtained at period $s - 1$ is evenly distributed in the network at time $k + 1$. We, therefore, introduce the *disagreement metric* ρ ,

$$\rho(k, s) = \max_{i, j \in \mathcal{M}} \left| [\Phi(k, s)]_{ij} - \frac{1}{m} \right| \quad \text{for all } k \geq s \geq 0, \tag{8}$$

which, when close to zero, establishes that all information obtained at time $s - 1$ by all agents is close to being evenly distributed in the network by time $k + 1$.

The analysis in the rest of this section is intended to produce upper bounds on the disagreement metric $\rho(k, s)$. We start our analysis by establishing an upper bound on the maximum distance between estimates of any two agents at any time k . In view of our communication model [cf. Eq. (6)], this bound will be essential in constructing positive probability events that ensure information gets propagated across the agents in the network.

Lemma 1 *Let Assumptions 1 and 3 hold. Let $x_i(k)$ be generated by the update rule in (2). Then, we have the following upper bound on the norm of the difference between*

the agent estimates: for all $k \geq 0$,

$$\max_{i,h \in \mathcal{M}} \|x_i(k) - x_h(k)\| \leq \Delta + 2mL \sum_{r=0}^{k-1} \alpha(r) + 2 \sum_{r=0}^{k-1} \sum_{j=1}^m \|e_j(r)\|,$$

where $\Delta = 2m \max_{j \in \mathcal{M}} \|x_j(0)\|$, and $e_j(k)$ denotes the projection error.

The lemma above establishes a bound on the distance between the agents' estimates that depends on the projection errors e_j , which are endogenously determined by the algorithm. However, if there exists some $M > 0$ such that $\|e_i(k)\| \leq M\alpha(k)$ for all $i \in \mathcal{M}$ and all $k \geq 0$, then the lemma above implies that, with probability 1, $\max_{i,h \in \mathcal{M}} \|x_i(k) - x_h(k)\| \leq \Delta + 2m(L + M) \sum_{r=0}^{k-1} \alpha(r)$. Under the assumption that such an M exists, we define the following set for each $k \in \mathbb{N}$,

$$R_M(k) = \left\{ x \in \mathbb{R}^{m \times n} \mid \max_{i,h \in \mathcal{M}} \|x_i(k) - x_h(k)\| \leq \Delta + 2m(L + M) \sum_{r=0}^{k-1} \alpha(r) \right\}. \tag{9}$$

This set represents the set of agent states which can be reached when the agents use the projected subgradient algorithm.

We next construct a sequence of events, denoted by $G(k)$, whose individual occurrence implies that information has been propagated from one agent to all other agents, therefore, implying a contraction of the disagreement metric ρ .

We say a link (j, i) is *activated at time k* when $a_{ij}(k) \geq \gamma$, and we denote by $\mathcal{E}(k)$ the set of such edges, i.e., $\mathcal{E}(k) = \{(j, i) \mid a_{ij}(k) \geq \gamma\}$. Here we construct an event in which the edges of the graphs $\mathcal{E}(k)$ are activated sequentially over time k , so that information propagates from every agent to every other agent in the network.

To define this event, we fix a node $w \in \mathcal{M}$ and consider *two directed spanning trees* rooted at w in the graph $(\mathcal{M}, \mathcal{E})$: an in-tree $T_{in,w}$ and an out-tree $T_{out,w}$. In $T_{in,w}$ there exists a directed path from every node $i \neq w$ to w ; while in $T_{out,w}$, there exists a directed path from w to every node $i \neq w$. The connectivity assumption imposed on $(\mathcal{M}, \mathcal{E})$ guarantees that these spanning trees exist and each contains $m - 1$ edges.

We order the edges of these spanning trees in a way such that on any directed path from a node $i \neq w$ to node w , edges are labeled in nondecreasing order. Let us represent the edges of the two spanning trees with the order described above as $T_{in,w} = \{e_1, e_2, \dots, e_{n-1}\}$ and $T_{out,w} = \{f_1, f_2, \dots, f_{n-1}\}$. For a detailed explanation of the process of building these spanning trees, see [14].

For all $l = 1, \dots, m - 1$, and any time $k \geq 0$, consider the events $B_l(k) = \{\omega \in \Omega \mid a_{e_l}(k + l - 1) \geq \gamma\}$ and $D_l(k) = \{\omega \in \Omega \mid a_{f_l}(k + (m - 1) + l - 1) \geq \gamma\}$, and define,

$$G(k) = \bigcap_{l=1}^{m-1} (B_l(k) \cap D_l(k)). \tag{10}$$

For all $l = 1, \dots, m - 1$, $B_l(k)$ denotes the event that edge $e_l \in T_{in,w}$ is activated at time $k + l - 1$, while $D_l(k)$ denotes the event that edge $f_l \in T_{out,w}$ is activated at

time $k + (m - 1) + l - 1$. Hence, $G(k)$ denotes the event in which each edge in the spanning trees $T_{in,w}$ and $T_{out,w}$ are activated sequentially following time k .

The following result establishes a bound on the probability of occurrence of such a $G(k)$ event. It states that the probability of an event $G(k)$ can be bounded as if the link activations were independent and each link activation had probability of occurring at least $\min \left\{ \delta, K / (\Delta + 2m(L + M) \sum_{r=1}^{k+2m-3} \alpha(r))^C \right\}$, where the $k + 2m - 3$ follows from the fact that event $G(k)$ is an intersection of $2(m - 1)$ events occurring consecutively starting at period k .

Lemma 2 *Let Assumptions 1, 2 and 3 hold. Let Δ denote the constant defined in Lemma 1. Moreover, assume that there exists $M > 0$ such that $\|e_i(k)\| \leq M\alpha(k)$ for all i and $k \geq 0$. Then,*

(a) *For all $s \in \mathbb{N}$, $k \geq s$, and any state $\bar{x} \in R_M(s)$,*

$$P(G(k)|x(s) = \bar{x}) \geq \min \left\{ \delta, \frac{K}{(\Delta + 2m(L + M) \sum_{r=1}^{k+2m-3} \alpha(r))^C} \right\}^{2(m-1)}.$$

(b) *For all $k \geq 0$, $u \geq 1$, and any state $\bar{x} \in R_M(k)$,*

$$\begin{aligned} &P \left(\bigcup_{l=0}^{u-1} G(k + 2(m - 1)l) \mid x(k) = \bar{x} \right) \\ &\geq 1 - \left(1 - \min \left\{ \delta, \frac{K}{(\Delta + 2m(L + M) \sum_{r=1}^{k+2(m-1)u-1} \alpha(r))^C} \right\}^{2(m-1)} \right)^u. \end{aligned}$$

The previous lemma bounded the probability of an event $G(k)$ occurring. The following lemma shows the implication of the event $G(k)$ for the disagreement metric.

Lemma 3 *Let Assumptions 2, 3 and 4 hold. Let t be a positive integer, and let there be scalars $s < s_1 < s_2 < \dots < s_t < k$, such that $s_{i+1} - s_i \geq 2(m - 1)$ for all $i = 1, \dots, t - 1$. For a fixed realization $\omega \in \Omega$, suppose that events $G(s_i)$ occur for each $i = 1, \dots, t$. Then, $\rho(k, s) \leq 2 \left(1 + \frac{1}{\gamma^{2(m-1)}} \right) (1 - \gamma^{2(m-1)})^t$.*

We skip the proof of this lemma since it would mirror the proof of Lemma 6 in [14]. We now obtain two propositions that establish contraction bounds on the disagreement metric based on two different sets of assumptions. For our first contraction bound, we need the following assumption on the sequence of stepsizes.

Assumption 5 (*Limiting stepsizes*) The sequence of stepsizes $\{\alpha(k)\}_{k \in \mathbb{N}}$ satisfies $\lim_{k \rightarrow \infty} k \log^p(k) \alpha(k) = 0$ for all $p < 1$.

The following lemma highlights two properties of stepsizes that satisfy Assumption 5: they are always square summable and they are not necessarily summable. The convergence results in Sect. 4 use stepsizes that are not summable, but square summable. An example of a stepsize that satisfies both of these properties, while also satisfying Assumption 5 is $\alpha(k) = 1/(k \log(k))$, for $k \geq 2$.

Lemma 4 Let $\{\alpha(k)\}_{k \in \mathbb{N}}$ be a stepsize sequence that satisfies Assumption 5. Then, the stepsizes are square summable, i.e., $\sum_{k=0}^{\infty} \alpha^2(k) < \infty$. Moreover, there exists a sequence of stepsizes $\{\bar{\alpha}(k)\}_{k \in \mathbb{N}}$ that satisfies Assumption 5 and is not summable, i.e., $\sum_{k=0}^{\infty} \bar{\alpha}(k) = \infty$.

The following proposition is one of the central results in our paper. It establishes, first, that for any fixed s , the expected disagreement metric $E[\rho(k, s) | x(s) = \bar{x}]$ decays at a rate of $e^{\sqrt{k-s}}$ as k goes to infinity. Importantly, it also establishes that, as s grows, the contraction bound for a fixed distance $k - s$ increases slowly in s . This slow increase is quantified by a function $\beta(s)$ that grows to infinity slower than the polynomial s^q for any $q > 0$.

Proposition 2 Let Assumptions 1, 2, 3, 4, and 5 hold. Assume also that there exists some $M > 0$ such that $\|e_i(k)\| \leq M\alpha(k)$ for all $i \in \mathcal{M}$ and $k \in \mathbb{N}$. Then, there exists a scalar $\mu > 0$, an increasing function $\beta(s) : \mathbb{N} \rightarrow \mathbb{R}_+$ and a function $S(q) : \mathbb{N} \rightarrow \mathbb{N}$ such that

$$\beta(s) \leq s^q \quad \text{for all } q > 0 \text{ and all } s \geq S(q) \tag{11}$$

$$\text{and } E[\rho(k, s) | x(s) = \bar{x}] \leq \beta(s)e^{-\mu\sqrt{k-s}} \quad \text{for all } k \geq s \geq 0, \bar{x} \in R_M(s). \tag{12}$$

Proof Part I. The first step of the proof is to define two functions, $g(k)$ and $w(k)$, that respectively bound the sum of the stepsizes up to time k and the inverse of the probability of communication at time k , and prove some limit properties of the functions $g(k)$ and $w(k)$ [see Eqs. (13) and (15)]. Define $g(k) : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ to be the linear interpolation of $\sum_{r=0}^{\lfloor k \rfloor} \alpha(k)$, i.e., $g(k) = \sum_{r=0}^{\lfloor k \rfloor} \alpha(k) + (k - \lfloor k \rfloor)\alpha(k - \lfloor k \rfloor + 1)$. Note that g is differentiable everywhere except at integer points and $g'(k) = \alpha(k - \lfloor k \rfloor + 1) = \alpha(\lceil k \rceil)$ at $k \notin \mathbb{N}$. We thus obtain from Assumption 5 that for all $p < 1$,

$$\lim_{k \rightarrow \infty, k \notin \mathbb{N}} k \log^p(k)g'(k) = \lim_{k \rightarrow \infty} \lceil k \rceil \log^p(\lceil k \rceil)\alpha(\lceil k \rceil) = 0. \tag{13}$$

Define $w(k)$ according to

$$w(k) = \frac{(\Delta + 2m(L + M)g(k))^{2(m-1)C}}{K^{2(m-1)}}, \tag{14}$$

where $\Delta = 2m \max_{j \in \mathcal{M}} \|x_j(0)\|$ and K and C are parameters of the communication model [see Eq. (6)]. We now show that for any $p < 1$,

$$\lim_{k \rightarrow \infty, k \notin \mathbb{N}} k \log^p(k)w'(k) = 0. \tag{15}$$

If $\lim_{k \rightarrow \infty} w(k) < \infty$, then the equation above holds immediately from Eq. (13). Therefore, assume $\lim_{k \rightarrow \infty} w(k) = \infty$. By L'Hospital's Rule, for any $q > 0$,

$$\lim_{k \rightarrow \infty, k \notin \mathbb{N}} \frac{w(k)}{\log^q(k)} = \frac{1}{q} \lim_{k \rightarrow \infty, k \notin \mathbb{N}} \frac{kw'(k)}{\log^{q-1}(k)}. \tag{16}$$

At the same time, if we take $w(k)$ to the power $\frac{1}{2(m-1)C}$ before using L'Hospital's Rule, we obtain that for any $q > 0$,

$$\begin{aligned} \lim_{k \rightarrow \infty, k \notin \mathbb{N}} \left(\frac{w(k)}{\log^q(k)} \right)^{\frac{1}{2(m-1)C}} &= \frac{1}{K^{1/C}} \lim_{k \rightarrow \infty, k \notin \mathbb{N}} \frac{\Delta + 2m(L + M)g(k)}{\log^{\frac{q}{2(m-1)C}}(k)} \\ &= \frac{4m(m-1)(L + M)C}{K^{1/C}q} \lim_{k \rightarrow \infty, k \notin \mathbb{N}} \frac{kg'(k)}{\log^{\frac{q}{2(m-1)C}-1}(k)} = 0, \end{aligned}$$

where the last equality follows from Eq. (13). From the equation above, we obtain

$$\lim_{k \rightarrow \infty, k \notin \mathbb{N}} \frac{w(k)}{\log^q(k)} = \left[\lim_{k \rightarrow \infty, k \notin \mathbb{N}} \left(\frac{w(k)}{\log^q(k)} \right)^{\frac{1}{2(m-1)C}} \right]^{2(m-1)C} = 0, \tag{17}$$

which combined with Eq. (16), yields the desired Eq. (15) for any $p = 1 - q < 1$.

Part 2. The second step of the proof involves defining a family of events $\{H_i(s)\}_{i,s}$ that occur with probability at least $\phi > 0$. We will later prove that an occurrence of $H_i(s)$ implies a contraction of the distance between the estimates. Let $h_i(s) = i + \lceil w(2s) \rceil$ for any $i, s \in \mathbb{N}$, where $w(\cdot)$ is defined in Eq. (14). We say the event $H_i(s)$ occurs if one out of a sequence of G -events [see definition in Eq. (10)] starting after s occurs. In particular, $H_i(s)$ is the union of $h_i(s)$ G -events and is defined as follows, $H_i(s) = \bigcup_{j=1}^{h_i(s)} G \left(s + 2(m-1) \left(j - 1 + \sum_{r=1}^{i-1} h_r(s) \right) \right)$ for all $i, s \in \mathbb{N}$, where $\sum_{r=1}^0(\cdot) = 0$. See Fig. 1 for a graphic representation of the $H_i(s)$ events.

We now show $P(H_i(s)|x(s) = \bar{x})$ for all $i, s \in \mathbb{N}$ and all $\bar{x} \in R_M(s)$ [see definition of $R_M(s)$ in Eq. (9)]. From Lemma 2(a) and the definition of $w(\cdot)$, we obtain that for all $\bar{x} \in R_M(s)$, $P(G(s)|x(s) = \bar{x}) \geq \min \left\{ \delta^{2(m-1)}, \frac{1}{w(s+2m-3)} \right\}$. Then, for all $s, i \in \mathbb{N}$ and all $\bar{x} \in R_M(s)$,

$$P(H_i(s)|x(s) = \bar{x}) \geq 1 - \left(1 - \min \left\{ \delta^{2(m-1)}, w \left(s + 2(m-1) \sum_{r=1}^i h_r(s) \right)^{-1} \right\} \right)^{h_i(s)},$$

where the inequality follows from Lemma 2(b) and the fact that $w(\cdot)$ is a non-decreasing function. Note that $h_r(s) \geq r$ for all r and s , so that $s + 2(m-1) \sum_{r=1}^i h_r(s) \geq i^2$. Let \hat{I} be the smallest i such that $w(i^2) \geq \delta^{-2(m-1)}$. We then have that for all $i \geq \hat{I}$, all s and all $\bar{x} \in R_M(s)$,

$$P(H_i(s)|x(s) = \bar{x}) \geq 1 - \left(1 - w \left(s + 2(m-1) \sum_{r=1}^i h_r(s) \right)^{-1} \right)^{h_i(s)},$$

Let \tilde{I} be the maximum between \hat{I} and the smallest i such that $w(i^2) > 1$. Using the inequality $(1 - 1/x)^x \leq e^{-1}$ for all $x \geq 1$, and multiplying and dividing the exponent in the equation above by $w \left(s + 2(m-1) \sum_{r=1}^i h_r(s) \right)$ we obtain $P(H_i(s)|x(s) = \bar{x})$




Fig. 1 The figure illustrates the three levels of probabilistic events considered in the proof: the events $B_l(s)$ and $D_l(s)$, which represent the occurrence of communication over a link (edge of the in-tree and out-tree, respectively); the events $G(s)$ as defined in (10), with length $2(m - 1)$ and whose occurrence dictates the spread of information from any agent to every other agent in the network; the events $H_i(s)$ constructed as the union of an increasing number of events $G(s)$ so that their probability of occurrence is guaranteed to be uniformly bounded away from zero. The occurrence of an event $H_i(s)$ also implies the spread of information from one agent to the entire network and, as a result, leads to a contraction of the distance between the agents' estimates

$\geq 1 - \exp\{-\frac{h_i(s)}{w(s+2(m-1)\sum_{r=1}^i h_r(s))}\}$ for all $i \geq \tilde{T}$, all s and all $\bar{x} \in R_M(s)$. By bounding $h_r(s) \leq h_i(s)$ and replacing $h_i(s) = i + \lceil w(2s) \rceil$, we obtain

$$P(H_i(s)|x(s) = \bar{x}) \geq 1 - e^{-\frac{i + \lceil w(2s) \rceil}{w(s+2(m-1)(i^2 + i\lceil w(2s) \rceil))}} \geq 1 - e^{-\frac{i + w(2s)}{w(s+2(m-1)(i^2 + iw(2s) + i))}}.$$

We now show there exists some \bar{T} such that

$$1 - e^{-\frac{i + w(2s)}{w(s+2(m-1)(i^2 + iw(2s) + i))}} \text{ is increasing in } i \text{ for all } i \geq \bar{T}, s \in \mathbb{N}. \tag{18}$$

The function above is increasing in i if $\frac{i + w(2s)}{w(s+2(m-1)(i^2 + iw(2s) + i))}$ is increasing in i . The partial derivative of this function with respect to i is positive if

$$w(s + 2(m - 1)(i^2 + iw(2s) + i)) - 2(m - 1)(2i^2 + i + 3iw(2s)) + w(2s) + w^2(2s)w'(s + 2(m - 1)(i^2 + iw(2s) + i)) > 0$$

at all points where the derivative $w'(\cdot)$ exists, that is, at non-integer values. If $i \geq \tilde{T}$, then $w(s + 2(m - 1)(i^2 + iw(2s) + i)) > 1$ for all s and it is thus sufficient to show

$$2(m - 1)(2i^2 + i + 3iw(2s) + w(2s) + w^2(2s))w'(s + 2(m - 1)(i^2 + iw(2s) + i)) \leq 1$$

in order to prove that Eq. (18) hold. The equation above holds if

$$2(m - 1)(3i^2 + 4iw(2s) + w^2(2s))w'(2(m - 1)(i^2 + iw(2s)) + s) \leq 1.$$

From Eq. (15) with $p = 1/2$, we have that there exists some N such that for all $x \geq N$, $w'(x) \leq \frac{1}{4x\sqrt{\log(x)}}$. For $i^2 \geq N$ and any $s \in \mathbb{N}$,

$$\begin{aligned} & 2(m - 1)(3i^2 + 4iw(2s) + w^2(2s))w'(2(m - 1)(i^2 + iw(2s)) + s) \\ & \leq \frac{2(m - 1)(3i^2 + 4iw(2s) + w^2(2s))}{(2(m - 1)(4i^2 + 4iw(2s)) + 4s)\sqrt{\log(2(m - 1)(i^2 + iw(2s)) + s)}} \\ & \leq \frac{3i^2 + 4iw(2s) + w^2(2s)}{4i^2 + 4iw(2s) + \frac{2}{m-1}s\sqrt{\log(i^2)}}. \end{aligned}$$

The term above is less than or equal to 1 if we select i large enough such that $\frac{2}{m-1}s\sqrt{\log(i^2)} \geq w^2(2s)$ for all $s \in \mathbb{N}$ [see Eq. (17) with $q < 1/2$], thus proving there exists some \bar{T} such that Eq. (18) holds. Hence, we obtain that for all $i, s \in \mathbb{N}$ and all $\bar{x} \in R_M(s)$, $P(H_i(s)|x(s) = \bar{x}) \geq$

$$\min_{j \in \{1, \dots, \bar{T}\}} \left\{ 1 - \left(1 - \min \left\{ \delta^{2(m-1)}, \frac{1}{w(s + 2(m - 1) \sum_{r=1}^j h_r(s))} \right\} \right)^{h_j(s)} \right\}.$$

Since $P(H_i(s)|x(s) = \bar{x}) > 0$ for all $i, s \in \mathbb{N}$ and all $\bar{x} \in R_M(s)$, to obtain the uniform lower bound on $P(H_i(s)|x(s) = \bar{x}) \geq \phi > 0$, it is sufficient to show that for all $i \in \{1, \dots, \bar{T}\}$ and all $\bar{x} \in R_M(s)$, $\lim_{s \rightarrow \infty} P(H_i(s)|x(s) = \bar{x}) > 0$. Repeating the steps above, but constraining s to be large enough instead of i , we obtain there exists some \tilde{S} such that for all $s \geq \tilde{S}$, all $i \in \mathbb{N}$ and $\bar{x} \in R_M(s)$, $P(H_i(s)|x(s) = \bar{x}) \geq 1 - \exp\{-\frac{i+w(2s)}{w(s+2(m-1)(i^2+iw(2s)+i))}\}$. Since there exists some \hat{S} such that $w(2s) \leq \log(2s)$ for all $s \geq \hat{S}$ [see Eq. (17) with $q = 1$], we obtain $P(H_i(s)|x(s) = \bar{x}) \geq 1 - \exp\{-\frac{i+w(2s)}{w(s+2(m-1)(i^2+i\log(2s)+i))}\}$ for $s \geq \max\{\hat{S}, \tilde{S}\}$ and all $i \in \mathbb{N}$ and $\bar{x} \in R_M(s)$.

Note that for every i , there exists some $\bar{S}(i)$ such that for all $s \geq \bar{S}(i)$ the numerator is greater than the denominator in the exponent above. Therefore, for all $i \in \mathbb{N}$ and $\bar{x} \in R_M(s)$, $\lim_{s \rightarrow \infty} P(H_i(s)|x(s) = \bar{x}) \geq 1 - e^{-1}$. Hence, there indeed exists some $\phi > 0$ such that $P(H_i(s)|x(s) = \bar{x}) \geq \phi$ for all $i, s \in \mathbb{N}$ and $\bar{x} \in R_M(s)$.

Part 3. In the previous step, we defined an event $H_i(s)$ and proved it had probability at least $\phi > 0$ of occurrence for any i and s . We now determine a lower bound on the number of possible H -events in an interval $\{s, \dots, k\}$. The maximum number of possible H events in the interval $\{s, \dots, k\}$ is given by $u(k, s) = \max\{t \in \mathbb{N} \mid s + 2(m - 1) \sum_{i=1}^t h_i(s) \leq k\}$. Recall that $h_i(s) = i + \lceil w(2s) \rceil \leq i + w(2s) + 1$ to obtain $u(k, s) \geq \max\{t \in \mathbb{N} \mid \sum_{i=1}^t (i + w(2s) + 1) \leq \frac{k-s}{2(m-1)}\}$. By expanding the sum and adding $(\frac{3}{2} + w(2s))^2$ to the left-hand side of the equation inside the maximization above, we obtain the following bound $u(k, s) \geq \max\{t \in \mathbb{N} \mid t^2 + 3t + 2w(2s)t + (\frac{3}{2} + w(2s))^2 \leq \frac{k-s}{m-1}\} = \max\{t \in \mathbb{N} \mid t + \frac{3}{2} + w(2s) \leq \sqrt{\frac{k-s}{m-1}}\}$, which yields the desired bound on $u(k, s)$,

$$u(k, s) \geq \sqrt{\frac{k-s}{m-1}} - \frac{5}{2} - w(2s). \tag{19}$$

Part 4. We now complete the proof of the proposition. The following argument shows there is a high probability that several H -events occur in a given $\{s, \dots, k\}$ interval and, therefore, we obtain the desired contraction.

Let $I_i(s)$ be the indicator variable of the event $H_i(s)$, that is $I_i(s) = 1$ if $H_i(s)$ occurs and $I_i(s) = 0$ otherwise. For any $k \geq s \geq 0$, any $\bar{x} \in R_M(s)$ and any $\delta > 0$, the disagreement metric ρ satisfies

$$\begin{aligned} & E[\rho(k, s) \mid x(s) = \bar{x}] \\ &= E \left[\rho(k, s) \mid x(s) = \bar{x}, \sum_{i=1}^{u(k,s)} I_i(s) > \delta u(k, s) \right] P \left(\sum_{i=1}^{u(k,s)} I_i(s) > \delta u(k, s) \mid x(s) = \bar{x} \right) \\ &+ E \left[\rho(k, s) \mid x(s) = \bar{x}, \sum_{i=1}^{u(k,s)} I_i(s) \leq \delta u(k, s) \right] P \left(\sum_{i=1}^{u(k,s)} I_i(s) \leq \delta u(k, s) \mid x(s) = \bar{x} \right). \end{aligned}$$

Since all the terms on the right-hand side of the equation above are less than or equal to 1, we obtain

$$\begin{aligned} & E[\rho(k, s) \mid x(s) = \bar{x}] \\ &\leq E \left[\rho(k, s) \mid x(s) = \bar{x}, \sum_{i=1}^{u(k,s)} I_i(s) > \delta u(k, s) \right] \\ &+ P \left(\sum_{i=1}^{u(k,s)} I_i(s) \leq \delta u(k, s) \mid x(s) = \bar{x} \right). \end{aligned} \tag{20}$$

We now bound the two terms in the right-hand side of Eq. (20). Consider initially the first term. If $I_i(s) > \delta u(k, s)$, then at least $\delta u(k, s)$ H -events occur, which by the definition of $H_i(s)$ implies that at least $\delta u(k, s)$ G -events occur. From Lemma 3, we obtain

$$\begin{aligned}
 E \left[\rho(k, s) \mid x(s) = \bar{x}, \sum_{i=1}^{u(k,s)} I_i(s) > \delta u(k, s) \right] \\
 \leq 2 \left(1 + \frac{1}{\gamma^{2(m-1)}} \right) \left(1 - \gamma^{2(m-1)} \right)^{\delta u(k,s)}
 \end{aligned} \tag{21}$$

for all $\delta > 0$. We now consider the second term in the right-hand side of Eq. (20). The events $\{I_i(s)\}_{i=1, \dots, u(k,s)}$ all have probability at least $\phi > 0$ conditional on any $x(s) \in R_M(s)$, but they are not independent. However, given any $x(s + \sum_{i=1}^{j-1} h_j(r)) \in R(s + \sum_{i=1}^{j-1} h_j(r))$, the event $I_j(s)$ is independent from the set of events $\{I_i(s)\}_{i=1, \dots, j-1}$ by the Markov property. Therefore, we can define a sequence of independent indicator variables $\{J_i(s)\}_{i=1, \dots, u(k,s)}$ such $P(J_i(s) = 1) = \phi$ and $J_i(s) \leq I_i(s)$ for all $i \in \{1, \dots, u(k, s)\}$ conditional on $x(s) \in R_M(s)$. Hence,

$$P \left(\sum_{i=1}^{u(k,s)} I_i(s) \leq \delta u(k, s) \mid x(s) = \bar{x} \right) \leq P \left(\sum_{i=1}^{u(k,s)} J_i(s) \leq \delta u(k, s) \right), \tag{22}$$

for any $\delta > 0$ and any $\bar{x} \in R_M(s)$. By selecting $\delta = \frac{\phi}{2}$ and using Hoeffding’s Inequality, we obtain

$$P \left(\frac{1}{u(k, s)} \sum_{i=1}^{u(k,s)} J_i(s) \leq \frac{\phi}{2} \right) \leq e^{-2\frac{\phi^2}{2^2}u(k,s)}. \tag{23}$$

Plugging Eqs. (21), (22) and (23), with $\delta = \phi/2$, into Eq. (20), we obtain $E[\rho(k, s)|x(s) = \bar{x}] \leq 2(1 + \frac{1}{\gamma^{2(m-1)}})(1 - \gamma^{2(m-1)})^{\frac{\phi}{2}u(k,s)} + e^{-\frac{\phi^2}{2}u(k,s)}$, for all $k \geq s \geq 0$ and all $\bar{x} \in R_M(s)$. This implies there exists some $\bar{\mu}_0, \bar{\mu}_1 > 0$ such that $\rho(k, s) \leq \bar{\mu}_0 e^{-\bar{\mu}_1 u(k,s)}$ and, combined with Eq. (19), we obtain there exist some $K, \mu > 0$ such that $E[\rho(k, s)|x(s) = \bar{x}] \leq K e^{\mu(w(2s) - \sqrt{k-s})}$ for all $k \geq s \geq 0, x(s) \in R_M(s)$. Let $\beta(s) = K e^{\mu w(2s)}$. Note that $\beta(\cdot)$ is an increasing function since $w(\cdot)$ is an increasing function. To complete the proof we need to show that β satisfies condition stipulated in Eq. (11). From Eq. (17), with $q = 1$, we obtain that $\lim_{s \rightarrow \infty, s \notin \mathbb{N}} \frac{w(s)}{\log(s)} = 0$. Note that since $w(\cdot)$ is a continuous function, the limit above also applies over the integers, i.e., $\lim_{s \rightarrow \infty} \frac{w(s)}{\log(s)} = 0$. Since $\lim_{s \rightarrow \infty} (s)$, for any $q > 0$, we have $0 = \lim_{s \rightarrow \infty} \frac{w(2s)}{\log(2s)} = \lim_{s \rightarrow \infty} \frac{\log(K) + \mu w(2s)}{q(-\log(2) + \log(2s))} = \lim_{s \rightarrow \infty} \frac{\log(\beta(s))}{\log(s^q)}$. Let $S(q)$ be a scalar such that $\frac{\log(\beta(s))}{\log(s^q)} \leq 1$ for all $s \geq S(q)$. We thus obtain that $\beta(s) \leq s^q$ for all $s \geq S(q)$, completing the proof of the proposition. \square

The above proposition yields the desired contraction of the disagreement metric ρ , but it assumes there exists some $M > 0$ such that $\|e_i(k)\| \leq M\alpha(k)$ for all $i \in \mathcal{M}$ and $k \in \mathbb{N}$. In settings where we do not have a guarantee that this assumption holds, we use the proposition below. Proposition 3 instead requires that the sets X_i be compact for each agent i . With compact feasible sets, the contraction bound on the disagreement

metric follows not from the prior analysis in this paper, but from an analysis of the information exchange that parallels the one we would use if the link activations were independent across time. This follows from the fact that the communication probability between agents $(j, i) \in \mathcal{E}$ is bounded by $\inf_{\bar{x} \in \prod_{i \in \mathcal{M}} X_i} \min\{\delta, K/\|\bar{x}_i - \bar{x}_j\|^C\}$, which is strictly positive when $\prod_{i \in \mathcal{M}} X_i$ is compact.

Proposition 3 *Let Assumptions 2, 3 and 4 hold. Assume also that the sets X_i are compact for all $i \in \mathcal{M}$. Then, there exist scalars $\kappa, \mu > 0$ such that for all $\bar{x} \in \prod_{i \in \mathcal{M}} X_i$,*

$$E[\rho(k, s)|x(s) = \bar{x}] \leq \kappa e^{-\mu(k-s)} \text{ for all } k \geq s \geq 0. \tag{24}$$

4 Analysis of the distributed subgradient method

In this section, we study the convergence behavior of the agent estimates $\{x_i(k)\}$ generated by the projected multi-agent subgradient algorithm (2). We first focus on the case when the constraint sets of agents are the same, i.e., for all $i, X_i = X$ for some closed convex nonempty set. In this case, we will prove almost sure consensus among agent estimates and almost sure convergence of agent estimates to an optimal solution when the stepsize sequence converges to 0 sufficiently fast (as stated in Assumption 5). We then consider the case when the constraint sets of the agents X_i are different convex compact sets and present convergence results both in terms of almost sure consensus of agent estimates and almost sure convergence of the agent estimates to an optimal solution under weaker assumptions on the stepsize sequence.

We first establish some key relations that hold under general stepsize rules that are used in the analysis of both cases.

4.1 Preliminary relations

The first relation measures the “distance” of the agent estimates to the intersection set $X = \cap_{i=1}^m X_i$. It will be key in studying the convergence behavior of the projection errors and the agent estimates. The properties of projection on a closed convex set, subgradients, and doubly stochasticity of agent weights play an important role in establishing this relation.

Lemma 5 *Let Assumption 3 hold. Let $\{x_i(k)\}$ and $\{e_i(k)\}$ be the sequences generated by the algorithm (3)–(5). For any $z \in X = \cap_{i=1}^m X_i$, the following hold:*

(a) *For all $k \geq 0$, we have*

$$\begin{aligned} \sum_{i=1}^m \|x_i(k+1) - z\|^2 &\leq \sum_{i=1}^m \|x_i(k) - z\|^2 + \alpha^2(k) \sum_{i=1}^m \|d_i(k)\|^2 \\ &\quad - 2\alpha(k) \sum_{i=1}^m (d_i(k))'(v_i(k) - z) - \sum_{i=1}^m \|e_i(k)\|^2. \end{aligned}$$

(b) *Let also Assumption 1 hold. For all $k \geq 0$, we have*

$$\begin{aligned} \sum_{i=1}^m \|x_i(k+1) - z\|^2 &\leq \sum_{i=1}^m \|x_i(k) - z\|^2 \\ &+ \alpha^2(k)mL^2 - 2\alpha(k) \sum_{i=1}^m (f_i(v_i(k)) - f_i(z)). \end{aligned} \tag{25}$$

Moreover, for all $k \geq 0$, it also follows that

$$\begin{aligned} \sum_{j=1}^m \|x_j(k+1) - z\|^2 &\leq \sum_{j=1}^m \|x_j(k) - z\|^2 + \alpha^2(k)mL^2 \\ &+ 2\alpha(k)L \sum_{j=1}^m \|x_j(k) - y(k)\| - 2\alpha(k) (f(y(k)) - f(z)), \end{aligned} \tag{26}$$

Our goal is to show that the agent disagreements $\|x_i(k) - x_j(k)\|$ converge to zero. To measure the agent disagreements $\|x_i(k) - x_j(k)\|$, we consider their average $\frac{1}{m} \sum_{j=1}^m x_j(k)$, and consider the disagreement of agent estimates with respect to this average. In particular, we define

$$y(k) = \frac{1}{m} \sum_{j=1}^m x_j(k) \quad \text{for all } k. \tag{27}$$

We have

$$y(k+1) = \frac{1}{m} \sum_{i=1}^m v_i(k) - \frac{\alpha(k)}{m} \sum_{i=1}^m d_i(k) + \frac{1}{m} \sum_{i=1}^m e_i(k).$$

When the weights are doubly stochastic, since $v_i(k) = \sum_{j=1}^m a_{ij}(k)x_j(k)$, it follows that

$$y(k+1) = y(k) - \frac{\alpha(k)}{m} \sum_{i=1}^m d_i(k) + \frac{1}{m} \sum_{i=1}^m e_i(k). \tag{28}$$

Under our assumptions, the next lemma provides an upper bound on the agent disagreements, measured by $\{\|x_i(k) - y(k)\|\}$ for all i , in terms of the subgradient bounds, projection errors and the disagreement metric $\rho(k, s)$ defined in Eq. (8).

Lemma 6 *Let Assumptions 1 and 3 hold. Let $\{x_i(k)\}$ be the sequence generated by the algorithm (3)–(5), and $\{y(k)\}$ be defined in Eq. (28). Then, for all i and $k \geq 2$, an upper bound on $\|x_i(k) - y(k)\|$ is given by*

$$\|x_i(k) - y(k)\| \leq m\rho(k - 1, 0) \sum_{j=1}^m \|x_j(0)\| + mL \sum_{r=0}^{k-2} \rho(k - 1, r + 1)\alpha(r) + 2\alpha(k - 1)L + \sum_{r=0}^{k-2} \rho(k - 1, r + 1) \sum_{j=1}^m \|e_j(r)\| + \|e_i(k - 1)\| + \frac{1}{m} \sum_{j=1}^m \|e_j(k - 1)\|.$$

In proving our convergence results, we will often use the following result on the infinite summability of products of positive scalar sequences with certain properties. This result was proven for geometric sequences in [19]. Here we use a generalized version for arbitrary summable sequences.

Lemma 7 *Let $\{\beta_l\}$ and $\{\gamma_k\}$ be positive scalar sequences, such that $\sum_{l=0}^\infty \beta_l < \infty$ and $\lim_{k \rightarrow \infty} \gamma_k = 0$. Then, $\lim_{k \rightarrow \infty} \sum_{\ell=0}^k \beta_{k-\ell} \gamma_\ell = 0$. In addition, if $\sum_{k=0}^\infty \gamma_k < \infty$, then $\sum_{k=0}^\infty \sum_{\ell=0}^k \beta_{k-\ell} \gamma_\ell < \infty$.*

4.2 Convergence analysis when $X_i = X$ for all i

In this subsection, we study the case when agent constraint sets X_i are the same. We study the asymptotic behavior of the agent estimates generated by the algorithm (2) using Assumption 5 on the stepsize sequence. The next assumption formalizes our condition on the constraint sets.

Assumption 6 *The constraint sets X_i are the same, i.e., $X_i = X$ for a closed convex set X .*

We show first that under this assumption, we can provide an upper bound on the norm of the projection error $\|e_i(k)\|$ as a function of the stepsize $\alpha(k)$ for all i and $k \geq 0$.

Lemma 8 *Let Assumptions 1 and 6 hold. Let $\{e_i(k)\}$ be the projection error defined by (5). Then, for all i and $k \geq 0$, the $e_i(k)$ satisfy $\|e_i(k)\| \leq 2L\alpha(k)$.*

This lemma shows that the projection errors are bounded by the scaled stepsize sequence under Assumption 6. Using this fact and an additional assumption on the stepsize sequence, we next show that the expected value of the sequences $\{\|x_i(k) - y(k)\|\}$ converge to zero for all i , thus establishing mean consensus among the agents in the limit. The proof relies on the bound on the expected disagreement metric $\rho(k, s)$ established in Proposition 2. The mean consensus result also immediately implies that the agent estimates reach almost sure consensus along a particular subsequence.

Proposition 4 *Let Assumptions 1, 2, 3, 4, and 6 hold. Assume also that the stepsize sequence $\{\alpha(k)\}$ satisfies Assumption 5. Let $\{x_i(k)\}$ be the sequence generated by the algorithm (3)–(5), and $\{y(k)\}$ be defined in Eq. (28). Then, for all i , we have*

$$\lim_{k \rightarrow \infty} E[\|x_i(k) - y(k)\|] = 0, \quad \text{and} \quad \liminf_{k \rightarrow \infty} \|x_i(k) - y(k)\| = 0 \quad \text{with probability one.}$$

The preceding proposition shows that the agent estimates reach a consensus in the expected sense. We next show that under Assumption 6, the agent estimates in fact converge to an almost sure consensus in the limit. We rely on the following standard convergence result for sequences of random variables, which is an immediate consequence of the supermartingale convergence theorem (see Bertsekas and Tsitsiklis [3]).

Lemma 9 Consider a probability space (Ω, F, P) and let $\{F(k)\}$ be an increasing sequence of σ -fields contained in F . Let $\{V(k)\}$ and $\{Z(k)\}$ be sequences of non-negative random variables (with finite expectation) adapted to $\{F(k)\}$ that satisfy $E[V(k + 1) | F(k)] \leq V(k) + Z(k)$ and $\sum_{k=1}^{\infty} E[Z(k)] < \infty$. Then, $V(k)$ converges with probability one, as $k \rightarrow \infty$.

Proposition 5 Let Assumptions 1, 2, 3, 4, and 6 hold. Assume also that the stepsize sequence $\{\alpha(k)\}$ satisfies Assumption 5. Let $\{x_i(k)\}$ be the sequence generated by the algorithm (3)–(5), and $\{y(k)\}$ be defined in Eq. (28). Then, for all i , we have:

- (a) $\sum_{k=2}^{\infty} \alpha(k) \|x_i(k) - y(k)\| < \infty$ with probability one.
- (b) $\lim_{k \rightarrow \infty} \|x_i(k) - y(k)\| = 0$ with probability one.

Proof (a) Using the upper bound on the projection error from Lemma 8, $\|e_i(k)\| \leq 2\alpha(k)L$ for all i and k , in Lemma 6, we have for all i and $k \geq 2$,

$$\begin{aligned} \|x_i(k) - y(k)\| &\leq m\rho(k - 1, 0) \sum_{j=1}^m \|x_j(0)\| \\ &\quad + 3mL \sum_{r=0}^{k-2} \rho(k - 1, r + 1)\alpha(r) + 6\alpha(k - 1)L. \end{aligned}$$

By multiplying this relation with $\alpha(k)$, we obtain

$$\begin{aligned} \alpha(k) \|x_i(k) - y(k)\| &\leq m\alpha(k)\rho(k - 1, 0) \sum_{j=1}^m \|x_j(0)\| \\ &\quad + 3mL \sum_{r=0}^{k-2} \rho(k - 1, r + 1)\alpha(k)\alpha(r) \\ &\quad + 6\alpha(k)\alpha(k - 1)L. \end{aligned}$$

Taking the expectation and using the estimate from Proposition 2, i.e., $E[\rho(k, s)] \leq \beta(s)e^{-\mu\sqrt{k-s}}$ for all $k \geq s \geq 0$, where μ is a positive scalar and $\beta(s)$ is a increasing sequence such that $\beta(s) \leq s^q$ for all $q > 0$ and all $s \geq S(q)$ for some integer $S(q)$, we have

$$\begin{aligned}
 E[\alpha(k)\|x_i(k) - y(k)\|] &\leq m\alpha(k)\beta(0)e^{-\mu\sqrt{k-1}} \sum_{j=1}^m \|x_j(0)\| \\
 &\quad + 3mL \sum_{r=0}^{k-2} \beta(r+1)e^{-\mu\sqrt{k-r-2}}\alpha(k)\alpha(r) \\
 &\quad + 6\alpha(k)\alpha(k-1)L.
 \end{aligned}$$

Let $\xi(r) = \beta(r+1)\alpha(r)$ for all $r \geq 0$. Using the relations $\alpha(k)\xi(r) \leq \alpha^2(k) + \xi^2(r)$ and $2\alpha(k)\alpha(k-1) \leq \alpha^2(k) + \alpha^2(k-1)$ for any k and r , the preceding implies that

$$\begin{aligned}
 E[\alpha(k)\|x_i(k) - y(k)\|] &\leq m\alpha(k)\beta(0)e^{-\mu\sqrt{k-1}} \sum_{j=1}^m \|x_j(0)\| \\
 &\quad + 3mL \sum_{r=0}^{k-2} e^{-\mu\sqrt{k-r-2}}\xi^2(r) \\
 &\quad + 3L\alpha^2(k) \left(m \sum_{r=0}^{k-2} e^{-\mu\sqrt{k-r-2}} + 1 \right) + 3\alpha^2(k-1)L.
 \end{aligned}$$

Summing over $k \geq 2$, we obtain

$$\begin{aligned}
 &\sum_{k=2}^{\infty} E[\alpha(k)\|x_i(k) - y(k)\|] \\
 &\leq m \sum_{j=1}^m \|x_j(0)\| \beta(0) \sum_{k=2}^{\infty} \alpha(k)e^{-\mu\sqrt{k-1}} \\
 &\quad + 3L \sum_{k=2}^{\infty} \left(\left(m \sum_{r=0}^{k-2} e^{-\mu\sqrt{k-r-2}} + 1 \right) \alpha^2(k) + \alpha^2(k-1) \right) \\
 &\quad + 3mL \sum_{k=2}^{\infty} \sum_{r=0}^{k-2} e^{-\mu\sqrt{k-r-2}}\xi^2(r).
 \end{aligned}$$

We next show the right handside of the above inequality is finite: since $\lim_{k \rightarrow \infty} \alpha(k) = 0$ (cf. Assumption 5), $\beta(0)$ is bounded, and $\sum_k e^{-\mu\sqrt{k}} < \infty$, Lemma 7 implies that the first term is bounded. The second term is bounded since $\sum_k \alpha^2(k) < \infty$ by Assumption 5 and Lemma 4. Since $\xi(r) = \beta(r+1)\alpha(r)$, we have for some small $\epsilon > 0$ and all r sufficiently large

$$\xi^2(r) = \beta^2(r+1)\alpha^2(r) \leq (r+1)^{2/3}\alpha^2(r) \leq (r+1)^{2/3} \frac{\epsilon}{r^2},$$

where the first inequality follows using the estimate in Eq. (11) with $q = 1/3$ and the second inequality follows from Assumption 5. This implies that $\sum_k \xi^2(k) < \infty$, which combined with Lemma 7 implies that the third term is also bounded. Hence, we have $\sum_{k=2}^\infty E[\alpha(k)\|x_i(k) - y(k)\|] < \infty$. By the monotone convergence theorem, this implies that $E[\sum_{k=2}^\infty \alpha(k)\|y(k) - x_i(k)\|] < \infty$, and therefore $\sum_{k=2}^\infty \alpha(k)\|y(k) - x_i(k)\| < \infty$ with probability 1, concluding the proof of this part.

(b) Using the iterations (4) and (28), we obtain for all $k \geq 1$ and i ,

$$\begin{aligned}
 y(k+1) - x_i(k+1) &= \left(y(k) - \sum_{j=1}^m a_{ij}(k)x_j(k) \right) \\
 &\quad - \alpha(k) \left(\frac{1}{m} \sum_{j=1}^m d_j(k) - d_i(k) \right) \\
 &\quad + \left(\frac{1}{m} \sum_{j=1}^m e_j(k) - e_i(k) \right).
 \end{aligned}$$

By the stochasticity of the weights $a_{ij}(k)$ and the subgradient boundedness, this implies that

$$\begin{aligned}
 &\|y(k+1) - x_i(k+1)\| \\
 &\leq \sum_{j=1}^m a_{ij}(k)\|y(k) - x_j(k)\| + 2L\alpha(k) + \frac{2}{m} \sum_{j=1}^m \|e_j(k)\|.
 \end{aligned}$$

Using the bound on the projection error from Lemma 5, we can simplify this relation as $\|y(k+1) - x_i(k+1)\| \leq \sum_{j=1}^m a_{ij}(k)\|y(k) - x_j(k)\| + 6L\alpha(k)$. Taking the square of both sides and using the convexity of the squared-norm function $\|\cdot\|^2$, this yields

$$\begin{aligned}
 \|y(k+1) - x_i(k+1)\|^2 &\leq \sum_{j=1}^m a_{ij}(k) \left[\|y(k) - x_j(k)\|^2 \right. \\
 &\quad \left. + 12L\alpha(k)\|y(k) - x_j(k)\| \right] + 36L^2\alpha(k)^2.
 \end{aligned}$$

Summing over all i and using the doubly stochasticity of the weights $a_{ij}(k)$, we have for all $k \geq 1$,

$$\begin{aligned}
 \sum_{i=1}^m \|y(k+1) - x_i(k+1)\|^2 &\leq \sum_{i=1}^m \|y(k) - x_i(k)\|^2 \\
 &\quad + 12L\alpha(k) \sum_{i=1}^m \|y(k) - x_i(k)\| + 36L^2m\alpha(k)^2.
 \end{aligned}$$

By part (a) of this lemma, we have $\sum_{k=1}^{\infty} \alpha(k) \|y(k) - x_i(k)\| < \infty$ with probability one. Since, we also have $\sum_k \alpha^2(k) < \infty$ (cf. Lemma 4), Lemma 9 applies and implies that $\sum_{i=1}^m \|y(k) - x_i(k)\|^2$ converges with probability one, as $k \rightarrow \infty$. By Proposition 4, we have $\liminf_{k \rightarrow \infty} \|x_i(k) - y(k)\| = 0$ with probability one. Since $\sum_{i=1}^m \|y(k) - x_i(k)\|^2$ converges with probability one, this implies that for all i , $\lim_{k \rightarrow \infty} \|x_i(k) - y(k)\| = 0$ with probability one, completing the proof. \square

We next present our main convergence result under Assumption 5 on the stepsize and Assumption 6 on the constraint sets.

Theorem 1 *Let Assumptions 1, 2, 3, 4 and 6 hold. Assume also that the stepsize sequence $\{\alpha(k)\}$ satisfies $\sum_{k=0}^{\infty} \alpha(k) = \infty$ and Assumption 5. Let $\{x_i(k)\}$ be the sequence generated by the algorithm (3)–(5). Then, there exists an optimal solution $x^* \in X^*$ such that for all i , $\lim_{k \rightarrow \infty} x_i(k) = x^*$ with probability one.*

Proof From Lemma 5(b), we have for some $z^* \in X^*$ (i.e., $f(z^*) = f^*$),

$$\begin{aligned} \sum_{j=1}^m \|x_j(k+1) - z^*\|^2 &\leq \sum_{j=1}^m \|x_j(k) - z^*\|^2 + \alpha^2(k)mL^2 \\ &\quad + 2\alpha(k)L \sum_{j=1}^m \|x_j(k) - y(k)\| - 2\alpha(k) (f(y(k)) - f^*), \end{aligned} \tag{29}$$

[see Eq. (26)]. Rearranging the terms and summing these relations over $k = 0, \dots, K$, we obtain

$$\begin{aligned} 2 \sum_{k=0}^K \alpha(k) (f(y(k)) - f^*) &\leq \sum_{j=1}^m \|x_j(0) - z^*\|^2 - \sum_{j=1}^m \|x_j(K+1) - z^*\|^2 \\ &\quad + mL^2 \sum_{k=0}^K \alpha^2(k) + 2L \sum_{k=0}^K \alpha(k) \sum_{j=1}^m \|x_j(k) - y(k)\|. \end{aligned}$$

By letting $K \rightarrow \infty$ in the relation above and using $\sum_{k=0}^{\infty} \alpha^2(k) < \infty$ (cf. Lemma 4) and $\sum_{k=0}^{\infty} \alpha(k) \sum_{j=1}^m \|x_j(k) - y(k)\| < \infty$ with probability one, we obtain that $\sum_{k=0}^K \alpha(k) (f(y(k)) - f^*) < \infty$ with probability one. Since $x_i(k) \in X$ for all i , we have $y(k) \in X$ [cf. Eq. (27)] and therefore $f(y(k)) \geq f^*$ for all k . Combined with the assumption $\sum_{k=0}^{\infty} \alpha(k) = \infty$, the preceding relation implies $\liminf_{k \rightarrow \infty} f(y(k)) = f^*$.

By dropping the nonnegative term $2\alpha(k) (f(y(k)) - f^*)$ in Eq. (29), we have

$$\sum_{j=1}^m \|x_j(k+1) - z^*\|^2 \leq \sum_{j=1}^m \|x_j(k) - z^*\|^2 + \alpha^2(k)mL^2 + 2\alpha(k)L \sum_{j=1}^m \|x_j(k) - y(k)\|.$$

Since $\sum_{k=0}^{\infty} \alpha^2(k) < \infty$ and $\sum_{k=0}^{\infty} \alpha(k) \sum_{j=1}^m \|x_j(k) - y(k)\| < \infty$ with probability one, Lemma 9 applies and implies that $\sum_{j=1}^m \|x_j(k) - z^*\|^2$ is a convergent sequence with probability one for all $z^* \in X^*$. By Lemma 5(b), we have $\lim_{k \rightarrow \infty} \|x_i(k) - y(k)\| = 0$ with probability one, therefore it also follows that the sequence $\|y(k) - z^*\|$ is also convergent. Since $y(k)$ is bounded, it must have a limit point. By the continuity of f (due to convexity of f over \mathbb{R}^n) and since $\liminf_{k \rightarrow \infty} f(y(k)) = f^*$, we obtain that one of the limit points of $\{y(k)\}$ must belong to X^* ; denote this limit point by x^* . Since the sequence $\{\|y(k) - x^*\|\}$ is convergent, it follows that $y(k)$ can have a unique limit point, i.e., $\lim_{k \rightarrow \infty} y(k) = x^*$ with probability one. This and $\lim_{k \rightarrow \infty} \|x_i(k) - y(k)\| = 0$ with probability one imply that each of the sequences $\{x_i(k)\}$ converges to the same $x^* \in X^*$ with probability one. \square

4.3 Convergence analysis for different constraint sets

In this subsection, we provide our convergence analysis for the case when all the constraint sets X_i are different. We show that even when the constraint sets of the agents are different, the agent estimates converge almost surely to an optimal solution of problem (1) under some conditions. In particular, we adopt the following assumption on the constraint sets.

Assumption 7 For each i , the constraint set X_i is convex and compact.

An important implication of the preceding assumption is that for each i , the subgradients of the function f_i at all points $x \in X_i$ are uniformly bounded, i.e., there exists some scalar $L > 0$ such that for all i , $\|d\| \leq L$ for all $d \in \partial f_i(x)$ and all $x \in X_i$.

Our first lemma shows that with different constraint sets and a stepsize that goes to zero, the projection error $e_i(k)$ converges to zero for all i along all sample paths.

Lemma 10 *Let Assumptions 3 and 7 hold. Let $\{x_i(k)\}$ and $\{e_i(k)\}$ be the sequences generated by the algorithm (3)–(5). Assume that the stepsize sequence satisfies $\alpha(k) \rightarrow 0$ as k goes to infinity.*

- (a) *For any $z \in X$, the scalar sequence $\sum_{i=1}^m \|x_i(k) - z\|^2$ is convergent.*
- (b) *The projection errors $e_i(k)$ converge to zero as $k \rightarrow \infty$, i.e., $\lim_{k \rightarrow \infty} \|e_i(k)\| = 0$ for all i .*

Proof (a) Using the subgradient boundedness and the relation $|d_i(k)'(v_i(k) - z)| \leq \|d_i(k)\| \|v_i(k) - z\|$ in part (a) of Lemma 5, we obtain

$$\begin{aligned} \sum_{i=1}^m \|x_i(k+1) - z\|^2 &\leq \sum_{i=1}^m \|x_i(k) - z\|^2 + \alpha^2(k)mL^2 \\ &\quad + 2\alpha(k)L \sum_{i=1}^m \|v_i(k) - z\| - \sum_{i=1}^m \|e_i(k)\|^2. \end{aligned}$$

Since $v_i(k) = \sum_{j=1}^m a_{ij}(k)x_j(k)$, using doubly stochasticity of the weights, we have $\sum_{i=1}^m \|v_i(k) - z\| \leq \sum_{i=1}^m \|x_i(k) - z\|$, which when combined with the preceding yields for any $z \in X$ and all $k \geq 0$,

$$\begin{aligned} \sum_{i=1}^m \|x_i(k + 1) - z\|^2 &\leq \sum_{i=1}^m \|x_i(k) - z\|^2 + \alpha^2(k)mL^2 \\ &\quad + 2\alpha(k)L \sum_{i=1}^m \|x_i(k) - z\| - \sum_{i=1}^m \|e_i(k)\|^2. \end{aligned} \tag{30}$$

Since $x_i(k) \in X_i$ for all i and X_i is compact (cf. Assumption 7), it follows that the sequence $\{x_i(k)\}$ is bounded for all i , and therefore the sequence $\sum_{i=1}^m \|x_i(k) - z\|$ is bounded. Since $\alpha(k) \rightarrow 0$ as $k \rightarrow \infty$, by dropping the nonnegative term $\sum_{i=1}^m \|e_i(k)\|^2$ in Eq. (30), it follows that

$$\begin{aligned} &\limsup_{k \rightarrow \infty} \sum_{i=1}^m \|x_i(k + 1) - z\|^2 \\ &\leq \liminf_{k \rightarrow \infty} \sum_{i=1}^m \|x_i(k) - z\|^2 \\ &\quad + \lim_{k \rightarrow \infty} \left(\alpha^2(k)mL^2 + 2\alpha(k)L \sum_{i=1}^m \|x_i(k) - z\| \right) \\ &= \liminf_{k \rightarrow \infty} \sum_{i=1}^m \|x_i(k) - z\|^2. \end{aligned}$$

Since the sequence $\sum_{i=1}^m \|x_i(k) - z\|^2$ is bounded, the preceding relation implies that the scalar sequence $\sum_{i=1}^m \|x_i(k) - z\|^2$ is convergent.

(b) From Eq. (30), for any $z \in X$, we have

$$\begin{aligned} \sum_{i=1}^m \|e_i(k)\|^2 &\leq \sum_{i=1}^m \|x_i(k) - z\|^2 - \sum_{i=1}^m \|x_i(k + 1) - z\|^2 + \alpha^2(k)mL^2 \\ &\quad + 2\alpha(k)L \sum_{i=1}^m \|x_i(k) - z\|. \end{aligned}$$

Taking the limit superior as $k \rightarrow \infty$, we obtain

$$\begin{aligned} \limsup_{k \rightarrow \infty} \sum_{i=1}^m \|e_i(k)\|^2 &\leq \lim_{k \rightarrow \infty} \left(\sum_{i=1}^m \|x_i(k) - z\|^2 - \sum_{i=1}^m \|x_i(k + 1) - z\|^2 \right) \\ &\quad + \lim_{k \rightarrow \infty} \left(\alpha^2(k)mL^2 + 2\alpha(k)L \sum_{i=1}^m \|x_i(k) - z\| \right), \end{aligned}$$

where the first term on the right handside is equal to zero by the convergence of the sequence $\sum_{i=1}^m \|x_i(k) - z\|^2$, and the second term is equal to zero by $\lim_{k \rightarrow \infty} \alpha(k) = 0$ and the boundedness of the sequence $\sum_{i=1}^m \|x_i(k) - z\|$, completing the proof. □

The preceding lemma shows the interesting result that the projection errors $\|e_i(k)\|$ converge to zero along all sample paths even when the agents have different constraint sets under the compactness conditions of Assumption 7. The next result establishes almost sure consensus among agent estimates.

Proposition 6 *Let Assumptions 2, 3, 4 and 7 hold. Let $\{x_i(k)\}$ be the sequence generated by the algorithm (3)–(5), and $\{y(k)\}$ be defined in Eq. (28). Assume that the stepsize sequence satisfies $\alpha(k) \rightarrow 0$. Then, for all i , we have $\lim_{k \rightarrow \infty} \|x_i(k) - y(k)\| = 0$ with probability one.*

The next theorem states our main convergence result for agent estimates when the constraint sets are different under some assumptions on the stepsize rule.

Theorem 2 *Let Assumptions 2, 3, 4 and 7 hold. Let $\{x_i(k)\}$ be the sequence generated by the algorithm (3)–(5). Assume that the stepsize sequence satisfies $\sum_k \alpha(k) = \infty$ and $\sum_k \alpha^2(k) < \infty$. Then, there exists an optimal solution $x^* \in X^*$ such that for all i $\lim_{k \rightarrow \infty} x_i(k) = x^*$ with probability one.*

Proof From Lemma 5(b), we have for some $z^* \in X^*$,

$$\begin{aligned} \sum_{i=1}^m \|x_i(k+1) - z^*\|^2 &\leq \sum_{i=1}^m \|x_i(k) - z^*\|^2 + \alpha^2(k) \sum_{i=1}^m \|d_i(k)\|^2 \\ &\quad - 2\alpha(k) \sum_{i=1}^m (f_i(v_i(k)) - f_i(z^*)). \end{aligned} \tag{31}$$

We show that the preceding implies that

$$\liminf_{k \rightarrow \infty} \sum_{i=1}^m f_i(v_i(k)) \leq f(z^*) = f^*. \tag{32}$$

Suppose to arrive at a contradiction that $\liminf_{k \rightarrow \infty} \sum_{i=1}^m f_i(v_i(k)) > f^*$. This implies that there exist some K and $\epsilon > 0$ such that for all $k \geq K$, we have $\sum_{i=1}^m f_i(v_i(k)) > f^* + \epsilon$. Summing the relation (31) over a window from K to N with $N > K$, we obtain

$$\sum_{i=1}^m \|x_i(N+1) - z^*\|^2 \leq \sum_{i=1}^m \|x_i(K) - z^*\|^2 + mL^2 \sum_{k=K}^N \alpha^2(k) - 2\epsilon \sum_{k=K}^N \alpha(k).$$

Letting $N \rightarrow \infty$, and using $\sum_k \alpha(k) = \infty$ and $\sum_k \alpha^2(k) < \infty$, this yields a contradiction and establishes the relation in Eq. (32).

By Proposition 6, we have

$$\lim_{k \rightarrow \infty} \|x_i(k) - y(k)\| = 0 \quad \text{with probability one.} \tag{33}$$

Since $v_i(k) = \sum_{j=1}^m a_{ij}(k)x_j(k)$, using the stochasticity of the weight vectors $a_i(k)$, this also implies

$$\lim_{k \rightarrow \infty} \|v_i(k) - y(k)\| \leq \lim_{k \rightarrow \infty} \sum_{j=1}^m a_{ij}(k) \|x_j(k) - y(k)\| = 0 \quad \text{with probability one.} \tag{34}$$

Combining Eqs. (32) and (34), we obtain $\liminf_{k \rightarrow \infty} f(y(k)) \leq f^*$ with probability one. From Lemma 10(a), the sequence $\{\sum_{i=1}^m \|x_i(k) - z^*\|\}$ is convergent for all $z^* \in X^*$. Combined with Eq. (33), this implies that the sequence $\{\|y(k) - z^*\|\}$ is convergent with probability one. Therefore, $y(k)$ is bounded and it must have a limit point. Moreover, since $x_i(k) \in X_i$ for all $k \geq 0$ and X_i is a closed set, all limit points of the sequence $\{x_i(k)\}$ must lie in the set X_i for all i . In view of Eq. (33), this implies that all limit points of the sequence $\{y(k)\}$ belong to the set X . Hence, from $\liminf_{k \rightarrow \infty} f(y(k)) \leq f^*$, we have $\liminf_{k \rightarrow \infty} f(y(k)) = f^*$ with probability one. Using the continuity of f (due to convexity of f over \mathbb{R}^n), this implies that one of the limit points of $\{y(k)\}$ must belong to X^* ; denote this limit point by x^* . Since the sequence $\{\|y(k) - x^*\|\}$ is convergent, it follows that $y(k)$ can have a unique limit point, i.e., $\lim_{k \rightarrow \infty} y(k) = x^*$ with probability one. This and $\lim_{k \rightarrow \infty} \|x_i(k) - y(k)\| = 0$ with probability one imply that each of the sequences $\{x_i(k)\}$ converges to the same $x^* \in X^*$ with probability one. \square

5 Conclusions

We studied distributed algorithms for multi-agent optimization problems over randomly-varying network topologies. We adopted a state-dependent communication model, in which the availability of links in the network is probabilistic with the probability dependent on the agent states. This is a good model for a variety of applications in which the state represents the position of the agents (in sensing and communication settings), or the beliefs of the agents (in social settings) and the distance of the agent states affects the communication and information exchange among the agents.

We studied a projected multi-agent subgradient algorithm for this problem and presented a convergence analysis for the agent estimates. The first step of our analysis establishes convergence rate bounds for a disagreement metric among the agent estimates. This bound is time-nonhomogeneous in that it depends on the initial time. Despite this, under the assumption that the stepsize sequence decreases sufficiently fast, we proved that agent estimates converge to an almost sure consensus and also to an optimal point of the global optimization problem under some assumptions on the constraint sets.

The framework introduced in this paper suggests a number of interesting further research directions. One future direction is to extend the constrained optimization problem to include both local and global constraints. This can be done using primal algorithms that involve projections, or using primal-dual algorithms where dual variables are used to ensure feasibility with respect to global constraints. Another interesting direction is to consider different probabilistic models for state-dependent communication. Our current model assumes the probability of communication is a continuous function of the l_2 norm of agent states. Considering other norms and discontinuous functions of agent states is an important extension which is relevant in a number of engineering and social settings.

Appendix A: notation and basic relations

A vector is viewed as a column vector, unless clearly stated otherwise. We denote by x_i or $[x]_i$ the i -th component of a vector x . When $x_i \geq 0$ for all components i of a vector x , we write $x \geq 0$. For a matrix A , we write A_{ij} or $[A]_{ij}$ to denote the matrix entry in the i -th row and j -th column. We write x' to denote the transpose of a vector x . The scalar product of two vectors $x, y \in \mathbb{R}^n$ is denoted by $x'y$. We use $\|x\|$ to denote the standard Euclidean norm, $\|x\| = \sqrt{x'x}$. We use the notion of a subgradient of a convex function $F(x)$ at a given vector $\bar{x} \in \text{dom}(F)$, the domain of F . We say that $s_F(\bar{x}) \in \mathbb{R}^n$ is a subgradient of the function F at $\bar{x} \in \text{dom}(F)$ when the following relation holds: $F(\bar{x}) + s_F(\bar{x})'(x - \bar{x}) \leq F(x)$ for all $x \in \text{dom}(F)$. The set of all subgradients of F at \bar{x} is denoted by $\partial F(\bar{x})$ (see [2]).

In our development, the properties of the projection operation on a closed convex set play an important role. Let X be a nonempty closed convex set in \mathbb{R}^n . We use $P_X[\bar{x}]$ to denote the projection of a vector \bar{x} on set X , i.e., $P_X[\bar{x}] = \arg \min_{x \in X} \|\bar{x} - x\|$. We will use the standard non-expansiveness property of projection, i.e.,

$$\|P_X[x] - P_X[y]\| \leq \|x - y\| \quad \text{for any } x \text{ and } y. \tag{35}$$

We will also use the following relation between the projection error vector and the feasible directions of the convex set X : for any $x \in \mathbb{R}^n$,

$$\|P_X[x] - y\|^2 \leq \|x - y\|^2 - \|P_X[x] - x\|^2 \quad \text{for all } y \in X. \tag{36}$$

References

1. Acemoglu, D., Ozdaglar, A., ParandehGheibi, A.: Spread of (mis)information in social networks. *Games Econ. Behav.* **70**, 194–227 (2010)
2. Bertsekas, D.P., Nedić, A., Ozdaglar, A.E.: *Convex analysis and optimization*. Athena Scientific, Cambridge, MA (2003)
3. Bertsekas, D.P., Tsitsiklis, J.N.: *Parallel and Distributed Computation: Numerical Methods*. Athena Scientific, Belmont, MA (1997)
4. Blondel, V.D., Hendrickx, J.M., Tsitsiklis, J.N.: On Krause’s multi-agent consensus model with state-dependent connectivity. *IEEE Trans. Automat. Control* **54**(11), 2586–2597 (2009)

5. Blondel, V.D., Hendrickx, J.M., Tsitsiklis, J.N.: Continuous-time average-preserving opinion dynamics with opinion-dependent communications. *SIAM J. Control Optim.* **48**(8), 5214–5240 (2010)
6. Blondel, V.D., Hendrickx, J.M., Olshevsky, A., Tsitsiklis, J.N.: Convergence in multiagent coordination, consensus, and flocking. In: *Proceedings of IEEE CDC* (2005)
7. Boyd, S., Ghosh, A., Prabhakar, B., Shah, D.: Gossip algorithms: design, analysis, and applications. In: *Proceedings of IEEE INFOCOM* (2005)
8. Cao, M., Spielman, D.A., Morse, A.S.: A lower bound on convergence of a distributed network consensus algorithm. In: *Proceedings of IEEE CDC* (2005)
9. Duchi, J., Agarwal, A., Wainwright, M.: Dual averaging for distributed optimization: Convergence analysis and network scaling, Preprint (2010)
10. Fagnani, F., Zampieri, S.: Randomized consensus algorithms over large scale networks. *IEEE J. Sel. Areas Commun.* **26**, 634–649 (2008)
11. Golub, B., Jackson, M.O.: Naive learning in social networks: Convergence, influence, and the wisdom of crowds. *Am. Econ. J. Microecon.* **2**(1), 112–149 (2010)
12. Hatano, Y., Mesbahi, M.: Agreement over random networks. *IEEE Trans. Automat. Control* **50**(11), 1867–1872 (2005)
13. Jadbabaie, A., Lin, J., Morse, S.: Coordination of groups of mobile autonomous agents using nearest neighbor rules. *IEEE Trans. Automat. Control* **48**(6), 988–1001 (2003)
14. Lobel, I., Ozdaglar, A.: Distributed subgradient methods for convex optimization over random networks. *IEEE Trans. Automat. Control* (2011) (to appear)
15. Lobel, I., Ozdaglar, A., Feijer, D.: Online appendices of distributed multi-agent optimization with state-dependent communication, <http://www.pages.stern.nyu.edu/~ilobel/state-dep-online-app.pdf> (2011)
16. Matei, I., Baras, J.: Distributed subgradient method under random communication topology—the effect of the probability distribution of the random graph on the performance metrics, ISR Technical Report 2009-15 (2009)
17. Nedić, A., Olshevsky, A., Ozdaglar, A., Tsitsiklis, J.N.: On distributed averaging algorithms and quantization effects. *IEEE Trans. Automat. Control* **54**(11), 2506–2517 (2009)
18. Nedić, A., Ozdaglar, A.: Distributed subgradient methods for multi-agent optimization. *IEEE Trans. Automat. Control* **54**(1), 48–61 (2009)
19. Nedić, A., Ozdaglar, A., Parrilo, P.A.: Constrained consensus and optimization in multi-agent networks. *IEEE Trans. Automat. Control* **55**, 922–938 (2010)
20. Nesterov, Y.: Primal-dual subgradient methods for convex problems. *Math. Program.* **120**(1), 261–283 (2009)
21. Olfati-Saber, R., Murray, R.M.: Consensus problems in networks of agents with switching topology and time-delays. *IEEE Trans. Automat. Control* **49**(9), 1520–1533 (2004)
22. Olshevsky, A., Tsitsiklis, J.N.: Convergence speed in distributed consensus and averaging. *SIAM J. Control Optim.* **48**(1), 33–55 (2009)
23. Ram, S.S., Nedić, A., Veeravalli, V.V.: Incremental stochastic subgradient algorithms for convex optimization. *SIAM J. Optim.* **20**(2), 691–717 (2009)
24. Tahbaz-Salehi, A., Jadbabaie, A.: A necessary and sufficient condition for consensus over random networks. *IEEE Trans. Automat. Control* **53**(3), 791–795 (2008)
25. Tsitsiklis, J.N., Bertsekas, D.P., Athans, M.: Distributed asynchronous deterministic and stochastic gradient optimization algorithms. *IEEE Trans. Automat. Control* **31**(9), 803–812 (1986)
26. Wu, C.W.: Synchronization and convergence of linear dynamics in random directed networks. *IEEE Trans. Automat. Control* **51**(7), 1207–1210 (2006)
27. Zhu, M., Martínez, S.: On distributed convex optimization under inequality and equality constraints via primal-dual subgradient methods, unpublished manuscript (2010)